

# THE CHARACTERISTIC POLYNOMIAL ON COMPACT GROUPS WITH HAAR MEASURE : SOME EQUALITIES IN LAW

P. BOURGADE, A. NIKEGHBALI, AND A. ROUAULT

Presented by Marc Yor.

ABSTRACT. This note presents some equalities in law for  $Z_N := \det(\text{Id} - G)$ , where  $G$  is an element of a subgroup of the set of unitary matrices of size  $N$ , endowed with its unique probability Haar measure. Indeed, under some general conditions,  $Z_N$  can be decomposed as a product of independent random variables, whose laws are explicitly known. Our results can be obtained in two ways : either by a recursive decomposition of the Haar measure (Section 1) or by previous results by Killip and Nenciu ([3]) on orthogonal polynomials with respect to some measure on the unit circle (Section 2). This latter method leads naturally to a study of determinants of a class of principal submatrices (Section 3).

RÉSUMÉ. Cette note présente quelques égalités en loi pour  $Z_N := \det(\text{Id} - G)$ , où  $G$  est un sous-groupe de l'ensemble des matrices unitaires de taille  $N$ , muni de son unique mesure de Haar normalisée. En effet, sous des conditions assez générales,  $Z_N$  peut être décomposé comme le produit de variables aléatoires indépendantes, dont on connaît la loi explicitement. Notre résultat peut être obtenu de deux manières: soit par une décomposition récursive de la mesure de Haar (Partie 1) soit en utilisant un résultat de Killip et Nenciu ([3]) à propos des polynômes orthogonaux relativement à une certaine mesure sur le cercle unité (Partie 2). Cette dernière méthode nous conduit naturellement à l'étude des déterminants de certaines sous-matrices (Partie 3).

In this note,  $\langle a, b \rangle$  denotes the Hermitian product of two elements  $a$  and  $b$  in  $\mathbb{C}^N$  (the dimension is implicit).

## 1. A RECURSIVE DECOMPOSITION, CONSEQUENCES

**1.1. The general equality in law.** Let  $\mathcal{G}$  be a subgroup of  $U(N)$ , the group of unitary matrices of size  $N$ . Let  $(e_1, \dots, e_N)$  be an orthonormal basis of  $\mathbb{C}^N$  and  $\mathcal{H} := \{H \in \mathcal{G} \mid H(e_1) = e_1\}$ , the subgroup of  $\mathcal{G}$  which stabilizes  $e_1$ . For a generic compact group  $\mathcal{A}$ , we write  $\mu_{\mathcal{A}}$  for the unique Haar probability measure on  $\mathcal{A}$ . Then we have the following Theorem.

**Theorem 1.1.** *Let  $M$  and  $H$  be independent matrices,  $M \in \mathcal{G}$  and  $H \in \mathcal{H}$  with distribution  $\mu_{\mathcal{H}}$ . Then  $MH \sim \mu_{\mathcal{G}}$  if and only if  $M(e_1) \sim f(\mu_{\mathcal{G}})$ , where  $f$  is the map  $f : G \mapsto G(e_1)$ .*

---

*Date:* 21 June 2007.

Let  $\mathcal{M}$  be the set of elements of  $\mathcal{G}$  which are reflections with respect to a hyperplane of  $\mathbb{C}^N$ . Define also

$$g : \begin{cases} \mathcal{H} & \rightarrow & U(N-1) \\ H & \mapsto & H_{\text{span}(e_2, \dots, e_N)} \end{cases},$$

where  $H_{\text{span}(e_2, \dots, e_N)}$  is the restriction of  $H$  to  $\text{span}(e_2, \dots, e_N)$ . Now suppose that  $\{G(e) \mid G \in \mathcal{G}\} = \{M(e) \mid M \in \mathcal{M}\}$ . Under this additional condition the following Theorem can be proven, using Theorem 1.1 and elementary manipulations of determinants.

**Theorem 1.2.** *Let  $G \sim \mu_G$ ,  $G' \sim \mu_{G'}$  and  $H \sim g(\mu_{\mathcal{H}})$  be independent. Then*

$$\det(\text{Id}_N - G) \stackrel{\text{law}}{=} (1 - \langle e_1, G'(e_1) \rangle) \det(\text{Id}_{N-1} - H).$$

**1.2. Examples : the unitary group, the group of permutations.** Take  $G = U(N)$ . As all reflections with respect to a hyperplane of  $\mathbb{C}^N$  are elements of  $G$ , iterations of Theorem 1.2 lead to the following Corollary.

**Corollary 1.3.** ([2]) *Let  $G \in U(N)$  be  $\mu_{U(N)}$  distributed. Then*

$$\det(\text{Id}_N - G) \stackrel{\text{law}}{=} \prod_{k=1}^N \left(1 - e^{i\theta_k} \sqrt{\beta_{1,k-1}}\right),$$

with  $\theta_1, \dots, \theta_N, \beta_{1,0}, \dots, \beta_{1,N-1}$  independent random variables, the  $\theta_k$ 's uniformly distributed on  $(0, 2\pi)$  and the  $\beta_{1,j}$ 's ( $0 \leq j \leq N-1$ ) being beta distributed with parameters 1 and  $j$  (by convention,  $\beta_{1,0}$  is the Dirac distribution at 1).

The group  $\mathcal{S}_N$  of permutations of size  $N$  gives another possible application. Identify an element  $\sigma \in \mathcal{S}_N$  with the matrix  $(\delta_{\sigma(i)}^j)_{1 \leq i, j \leq N}$  ( $\delta$  is Kronecker's symbol). As  $\det(\text{Id}_N - \sigma)$  is equal to 0, we prefer to deal with the group  $\tilde{\mathcal{S}}_N$  of matrices  $(e^{i\theta_j} \delta_{\sigma(i)}^j)_{1 \leq i, j \leq N}$ , with  $\sigma \in \mathcal{S}_N$  and  $\theta_1, \dots, \theta_N$  independent uniform random variables on  $(0, 2\pi)$ . As previously, iterations of Theorem 1.2 give the following result.

**Corollary 1.4.** *Let  $S_N \in \tilde{\mathcal{S}}_N$  be  $\mu_{\tilde{\mathcal{S}}_N}$  distributed. Then*

$$\det(\text{Id}_N - S_N) \stackrel{\text{law}}{=} \prod_{k=1}^N \left(1 - e^{i\theta_k} X_k\right),$$

with  $\theta_1, \dots, \theta_N, X_1, \dots, X_N$  independent random variables, the  $\theta_k$ 's uniformly distributed on  $(0, 2\pi)$  and the  $X_k$ 's Bernoulli variables :  $\mathbb{P}(X_k = 1) = 1/k$ ,  $\mathbb{P}(X_k = 0) = 1 - 1/k$ .

## 2. CHARACTERISTIC POLYNOMIALS AS ORTHOGONAL POLYNOMIALS

We now show how Corollary 1.3 can be obtained as a consequence of a result by Killip and Nenciu ([3]).

**2.1. A result by Killip and Nenciu.** Let  $\mathbb{D}$  be the open unit disk  $\{z \in \mathbb{C} : |z| < 1\}$  and  $\partial\mathbb{D}$  the unit circle. Let  $(e_1, \dots, e_N)$  be the canonical basis of  $\mathbb{C}^N$ . If  $G \in U(N)$ , and if  $e_1$  is cyclic for  $G$ , the spectral measure for the pair  $(G, e_1)$  is the unique probability  $\nu$  on  $\partial\mathbb{D}$  such that, for every integer  $k \geq 0$

$$\langle e_1, G^k e_1 \rangle = \int_{\partial\mathbb{D}} z^k d\nu(z). \quad (2.1)$$

In fact, we have the expression

$$\nu = \sum_{j=1}^N \pi_j \delta_{e^{i\zeta_j}}$$

where  $(e^{i\zeta_j}, j = 1, \dots, N)$  are the eigenvalues of  $G$  and where  $\pi_j = |\langle e_1, \Pi e_j \rangle|^2$  with  $\Pi$  a unitary matrix diagonalizing  $G$ .

The relation (2.1) allows to define an isometry from  $\mathbb{C}^N$  equipped with the basis  $(e_1, Ge_1, \dots, G^{N-1}e_1)$  into the subspace of  $L^2(\partial\mathbb{D}; d\nu)$  spanned by the family  $(1, z, \dots, z^N)$ . The endomorphism  $G$  is then a representation of the multiplication by  $z$ .

From the linearly independent family of monomials  $\{1, z, z^2, \dots, z^{N-1}\}$  in  $L^2(\partial\mathbb{D}, \nu)$ , we construct an orthogonal basis  $\Phi_0, \dots, \Phi_{N-1}$  of monic polynomials by the Gram-Schmidt procedure. The  $N^{\text{th}}$  degree polynomial obtained this way is

$$\Phi_N(z) = \prod_{j=1}^N (z - e^{i\zeta_j}),$$

i.e. the characteristic polynomial of  $G$ . The  $\Phi_k$ 's ( $k = 0, \dots, N$ ) obey the Szegő recursion relation:

$$\Phi_{j+1}(z) = z\Phi_j(z) - \bar{\alpha}_j \Phi_j^*(z) \quad (2.2)$$

where  $\Phi_j^*(z) = z^j \overline{\Phi_j(\bar{z}^{-1})}$ . The coefficients  $\alpha'_j$ s ( $j \geq 0$ ) are called Schur or Verblunsky coefficients and satisfy the condition  $\alpha_0, \dots, \alpha_{N-2} \in \mathbb{D}$  and  $\alpha_{N-1} \in \partial\mathbb{D}$ . There is a bijection between this set of coefficients and the set of spectral probability measures  $\nu$  (Verblunsky's theorem). If  $G \sim \mu_{U(N)}$ , then we know the exact distribution of the Verblunsky coefficients :

**Theorem 2.1.** (Killip and Nenciu [3]) *Let  $G \in U(N)$  be  $\mu_{U(N)}$  distributed. The Verblunsky parameters  $\alpha_0, \dots, \alpha_{N-2}, \alpha_{N-1}$  are independent and the density of  $\alpha_j$  for  $j \leq N - 1$  is*

$$\frac{N-j-1}{\pi} (1-|z|^2)^{N-j-2} \mathbf{1}_{\mathbb{D}}(z)$$

(for  $j = N - 1$  by convention this is the uniform measure on the unit circle).

**2.2. Recovering Corollary 1.3.** For  $z = 1$ , Szegő's recursion (2.2) can be written

$$\Phi_{j+1}(1) = \Phi_j(1) - \bar{\alpha}_j \overline{\Phi_j(1)}. \quad (2.3)$$

Under the Haar measure for  $G$ , as  $\alpha_j$  is independent of  $\Phi_j(1)$  and its distribution is invariant by rotation, (2.3) easily yields

$$\Phi_{j+1}(1) \stackrel{\text{law}}{=} (1 - \alpha_j) \Phi_j(1).$$

In particular, for  $j = N - 1$  we get by induction exactly the same result as Corollary 1.3.

*Remark.* A similar result holds for  $SO(2N)$ , and can be shown using either the method of Section 1 or the one in Section 2, with the corresponding result by Killip and Nenciu for the Verblunsky coefficients on the orthogonal group [3].

**2.3. Extension.** We now consider the whole sequence of polynomials  $\Phi_j, j \leq N$  for  $j \leq N$  as a sequence of characteristic polynomials. For this purpose, we apply the Gram-Schmidt procedure to  $1, z, z^{-1}, z^2, \dots, z^{p-1}, z^{1-p}, z^p$  if  $N = 2p$  and to  $1, z, z^{-1}, z^2, \dots, z^p, z^{-p}$  if  $N = 2p + 1$  in  $L^2(\partial\mathbb{D}); d\nu$ . In the resulting basis, the mapping  $f(z) \mapsto zf(z)$  is represented by a so-called CMV matrix ([3] Appendix B, [5]) denoted by  $\mathcal{C}_N(G)$ . It is five-diagonal and conjugate to  $G$ . For  $1 \leq j \leq N$  let  $\mathcal{C}_N^{(j)}(G)$  the principal submatrix of order  $j$  of  $\mathcal{C}_N(G)$ . It is known (see for instance Proposition 3.1 in [5]) that

$$\Phi_j(z) = \det \left( z\text{Id}_j - \mathcal{C}_N^{(j)}(G) \right).$$

From the recursion (2.3) and looking at the invariance of conditional distributions, we see that

$$\left( \det \left( \text{Id}_j - \mathcal{C}_N^{(j)}(G) \right) \right)_{1 \leq j \leq N} = (\Phi_j(1))_{1 \leq j \leq N} \stackrel{\text{law}}{=} \left( \prod_{l=0}^j (1 - \alpha_l) \right)_{0 \leq j \leq N-1}. \quad (2.4)$$

It allows a study of the process  $(\log \Phi_{\lfloor Nt \rfloor}(1))$ ,  $t \in [0, 1]$  as a triangular array of (complex) independent random variables. For  $t = 1$  the asymptotic behavior is presented in [2] (see (2.6 below). It is remarkable that for  $t < 1$ , we do not need any normalization for the CLT.

**Theorem 2.2.** (1) *As  $N \rightarrow \infty$*

$$\left( \log \det \left( \text{Id}_j - \mathcal{C}_{\lfloor Nt \rfloor}^{(j)}(G) \right); t \in [0, 1] \right) \Rightarrow \left( \mathbf{B}_{-\frac{1}{2} \log(1-t)}; t \in [0, 1] \right), \quad (2.5)$$

where  $\mathbf{B}$  is a standard complex Brownian motion and  $\Rightarrow$  stands for the weak convergence of distributions in the set of càdlàg functions on  $[0, 1]$ , starting from 0, endowed with the Skorokhod topology.

(2) *As  $N \rightarrow \infty$ ,*

$$\frac{\log \det(\text{Id}_N - G)}{\sqrt{2 \log N}} \Rightarrow \mathcal{N}_1 + i\mathcal{N}_2 \quad (2.6)$$

where  $\mathcal{N}_1$  and  $\mathcal{N}_2$  are independent standard normal and independent of  $\mathbf{B}$ , and  $\Rightarrow$  stands for the weak convergence of distributions in  $\mathbb{C}$ .

This theorem can be proved using the Mellin-Fourier transform of the  $1 - \alpha_j$ 's and independence. This method may also be used to prove large deviations. It is the topic of a companion paper. These results occur in similar way for other random determinants (see [4]).

## REFERENCES

- [1] R. Arratia, A.D. Barbour, S. Tavaré, Logarithmic Combinatorial Structures: A Probabilistic Approach. 352 pp, 2003. EMS Monographs in Mathematics, 1. European Mathematical Society Publishing House, Zürich.
- [2] P. Bourgade, C.P. Hughes, A. Nikeghbali, M. Yor, The characteristic polynomial of a random unitary matrix: a probabilistic approach, arXiv math.PR/0706.0333, 2007.
- [3] R. Killip and I. Nenciu, Matrix models for circular ensembles, International Mathematics Research Notices, vol. 2004, no. 50, pp. 2665-2701, 2004.
- [4] A. Rouault, Asymptotic behavior of random determinants in the Laguerre, Gram and Jacobi ensembles, arXiv math.PR/0607767, 2007.
- [5] B. Simon, CMV matrices: Five years after, arXiv math.SP/0603093, 2006.

LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES, UNIVERSITÉ PIERRE ET MARIE CURIE,  
ET C.N.R.S. UMR 7599, 175, RUE DU CHEVALERET, F-75013 PARIS, FRANCE  
*E-mail address:* bourgade@enst.fr

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, CH-8057 ZÜRICH,  
SWITZERLAND  
*E-mail address:* ashkan.nikeghbali@math.unizh.ch

UNIVERSITÉ VERSAILLES-SAINT QUENTIN, LMV, BÂTIMENT FERMAT, 45 AVENUE DES ETATS-UNIS,  
78035 VERSAILLES CEDEX  
*E-mail address:* rouault@fermat.math.uvsq.fr