

LARGE DEVIATIONS AND BRANCHING PROCESSES.

Alain ROUAULT, (Versailles, France)

ABSTRACT. These lecture notes are devoted to present several uses of Large Deviation asymptotics in Branching Processes.

Overview

I) Introduction .

Why large deviations ?

II) Background on Large Deviations

2.1 Cramer-Chernov theorem. Sums of i.i.d. r.v..

2.2 Large Deviation Principle.

2.3 Gärtner-Ellis theorem.

2.4 Varadhan theorem. Integration of functions.

2.5 Schilder theorem.

III) Branching processes. Notations and Models

3.1 Trees.

3.2 Marked trees.

IV) Large deviations and Galton-Watson processes

4.1 Asymptotical behaviour of generating function.

4.2 Growth rates.

4.3 Tail behaviour.

V) Large deviations in spatial processes

5.1 Rates of propagation

5.1.1 Branching Random Walk

5.1.2 Malthusian phenomena.

5.2 Growth rates in the supercritical area (Branching Random Walk).

5.3 Sharp asymptotics in the subcritical area

5.3.1 Branching Random Walk

5.3.2 Homogeneous Branching Brownian Motion

5.3.3 Inhomogeneous Branching Brownian Motion.

1. INTRODUCTION

The large deviation techniques are relevant in various topics of the branching processes theory. There are many intuitive reasons for that. Let us just give one.

As supercritical branching processes are connected with exponential growing of populations, and subcritical ones with exponential decreasing of presence probabilities, it is rather natural to find large deviations exponential estimates. In various supercritical spatial branching models, there are two phenomena : the branching structure and the process along a branch. If the latter one is a random walk, then the large deviation asymptotics for

sums of i.i.d. random variables plays an important role. At level n and speed a there is a competition between the exponentially small probability $\exp -nh(a)$ (where h is the Cramer transform of the motion) and the exponential number of branches m^n . If $h(a) - \log m < 0$ the effective of branches following approximately the speed a is roughly $\exp -n(h(a) - \log m)$. If $h(a) - \log m > 0$ then the probability of finding a branch of speed approximately a is very small, actually $\exp -n(h(a) - \log m)$. This topic is widely developed in Section 5 with two types of results, determination of the propagation front and precise results for populations (range of supercriticality) or probabilities (range of subcriticality).

In section 2, we give a background on large deviations as it is presented now in the literature (see for instance Dembo-Zeitouni), limiting ourselves to the main theorems. In section 3 we recall the notations and various models of branching processes. In section 4 we present some recent results of large deviations in Galton-Watson process (tail behavior of the limit r.v. W and growth rates of the generation size).

An excellent bibliography on Branching Processes is in Vatutin (1993). The survey of large deviations in branching processes given here is by no means complete. I omitted superprocesses by lack of space (time). The density dependent branching processes generate also nice large deviation theorems (Klebaner and Zeitouni (1994), Pierre-Loti-Viaud and Portal (1993)).

2. BACKGROUND ON LARGE DEVIATIONS

In this survey we state the main results with different notations, which allows to apply them directly in the subsequent paragraphs.

2.1. CRAMER-CHEBNOV THEOREM

Let μ be a probability on \mathbb{R} . The log-Laplace of μ defined from \mathbb{R} to $[0, \infty]$ by

$$\Lambda_\mu(\theta) = \log \int \exp(\theta x) \mu(dx),$$

is a convex function (apply Hölder inequality). Let Λ_μ^* be the Fenchel-Legendre dual of Λ_μ defined for $x \in \mathbb{R}$ by :

$$\Lambda_\mu^*(x) = \sup\{\theta x - \Lambda_\mu(\theta) : \theta \in \mathbb{R}\} \in [0, +\infty].$$

As a supremum of linear functions, Λ_μ^* is lower semi-continuous (l.s.c.) and convex. It is positive (take $\theta = 0$). The main feature of the following Cramer-Chernov theorem is that it needs no moment condition.

THEOREM 2.1.

Let μ a distribution on \mathbb{R} and for $n \geq 1$, let μ_n the distribution under $\mu^{\otimes n}$ of $\frac{1}{n} \sum_1^n x_i$.

a) For every closed $F \subset \mathbb{R}$,

$$\overline{\lim} \frac{1}{n} \log(\mu_n(F)) \leq -\inf\{\Lambda_\mu^*(x); x \in F\} \quad (2.1)$$

b) For every open $G \subset \mathbb{R}$,

$$\underline{\lim} \frac{1}{n} \log(\mu_n(G)) \geq -\inf\{\Lambda_\mu^*(x); x \in G\} \quad (2.2)$$

REMARK 2.2. Actually if μ have an expectation \underline{x} then we have a Cramer-Chernov upper bound holding for every n :

$$\mu_n(F) \leq 2 \exp -n \inf\{\Lambda_\mu^*(x); x \in F\} \quad (2.3)$$

If we assume that Λ_μ is defined everywhere, then Λ et Λ^* have some nice properties described in the following proposition. Let $[a, b]$ be the convex hull of the support of μ .

PROPOSITION 2.3. Under the above conditions,

- a) $\Lambda \in \mathcal{C}^\infty$.
- b) If X is a r.v. μ distributed, then $EX = \Lambda'(0)$, $Var X = \Lambda''(0)$.
- c) Λ is strictly convex .
- d) $\Lambda^* \geq 0$ with equality only in EX .
- e) $(\Lambda)'$ is one-to-one from \mathbb{R} onto $]a, b[$.
- f) $]a, b[\subseteq D_{\Lambda^*} \subseteq [a, b]$.
- g) If $z \in \{a, b\}$ then $z \in D_{\Lambda^*}$ if and only if $\mu(\{z\}) > 0$; in that case $\Lambda^*(z) = -\log \mu(\{z\})$. In particular, if $Supp \mu$ is a finite set then $D_{\Lambda^*} = [a, b]$ and Λ^* is continuous on its domain.
- h) Λ^* is differentiable on $]a, b[$ and $|(\Lambda^*)'(z)| \rightarrow \infty$ as $z \rightarrow a$ or b staying in $]a, b[$.
- i) $(\Lambda^*)'$ is one-to-one from $]a, b[$ onto \mathbb{R} and its inverse is $(\Lambda)'$.
- j) Λ^* is analytic on $]a, b[$.

2.2. LARGE DEVIATION PRINCIPLE

Let (E, \mathcal{E}) be a measurable space provided with a topology. A large deviation principle (LDP) describes the asymptotic behaviour, as $\epsilon \rightarrow 0$, of a family of probability measures (μ_ϵ) on (E, \mathcal{E}) by the means of a so-called rate function.

A function f from E to $[0, +\infty]$ is l.s.c. if for all $\alpha \in [0, +\infty[$ the level set $\Psi_f(\alpha) := \{x : f(x) \leq \alpha\}$ is closed in E .

DEFINITION 2.4. A rate function I is a l.s.c. mapping from E to $[0, +\infty]$. A good rate function is a rate function with compact level sets.

Recall that a l.s.c. function reaches its inf on compact sets. If it is good, it reaches its inf on closed sets.

DEFINITION 2.5. The family (μ_ϵ) satisfies a LDP of rate function I if

a) For every closed set $F \in \mathcal{E}$,

$$\limsup \epsilon \log \mu_\epsilon(F) \leq -\inf\{I(x); x \in F\}$$

b) For every open set $G \in \mathcal{E}$,

$$\liminf_\epsilon \epsilon \log \mu_\epsilon(G) \geq -\inf\{I(x); x \in G\}.$$

Since μ_ϵ is a probability, it is needed that $\inf\{I(x); x \in E\} = 0$ due to the upper bound. When I is good, $I = 0$ has at least one solution.

REMARK. The rate function associated with a LDP satisfied by (μ_ϵ) on a metric space is unique.

2.3. GÄRTNER-ELLIS THEOREM

We want to weaken assumptions about independence and dimension.

Let (Y_n) be a sequence of real r.v. defined on $\{(\Omega_n, \mathcal{F}_n, P_n); n = 1, 2, \dots\}$ and let (a_n) a sequence of positive numbers tending to infinity. Let

$$\Lambda_n(\theta) := \frac{1}{a_n} \log E_n\{\exp \langle \theta, Y_n \rangle\}$$

ASSUMPTION GE. For all $\theta \in \mathbb{R}^d$, the limit of $\Lambda_n(\theta)$, denoted $\Lambda(\theta)$ exists in $[-\infty, +\infty]$. Moreover $0 \in \text{int } D_\Lambda$.

For instance if $a_n = n$ and $Y_n = S_n$, with S_n a sum of i.i.d. r.v. we have $\Lambda_n \equiv \Lambda$.

Under GE, it is easy to see that Λ take its values in $]-\infty, +\infty]$ and is convex. Its dual Λ^* is l.s.c. and convex. Since $0 \in \text{int } D_\Lambda$ it is good and $\inf\{\Lambda^*(x); x \in \mathbb{R}^d\} = 0$.

DEFINITION 2.6. $y \in \mathbb{R}^d$ is an exposed point of Λ^* if there exists some $\theta \in \mathbb{R}^d$ such that for all $x \neq y$ we have

$$\langle \theta, y \rangle - \Lambda^*(y) > \langle \theta, x \rangle - \Lambda^*(x).$$

Such a θ is called exposed hyperplane, it belongs to the subdifferential of Λ^* in y .

DEFINITION 2.7. A convex function $\Lambda : \mathbb{R}^d \rightarrow]-\infty, +\infty]$ is called essentially smooth if

- a) $\text{int } D_\Lambda \neq \emptyset$
- b) Λ is differentiable on $\text{int } D_\Lambda$
- c) Λ is steep i.e. $\lim |\nabla \Lambda(\theta_n)| = +\infty$ for all sequence $\theta_n \in \text{int } D_\Lambda$ tending to a border point of $\text{int } D_\Lambda$.

THEOREM 2.8. Assume (GE). Let μ_n be the distribution of $\frac{Y_n}{a_n}$ sur \mathbb{R}^d ,

a) For all closed set $F \in \mathbb{R}^d$,

$$\overline{\lim} \frac{1}{a_n} \log \mu_n(F) \leq -\inf\{\Lambda^*(x); x \in F\}$$

b) For all open set $G \in \mathbb{R}^d$,

$$\underline{\lim} \frac{1}{a_n} \log \mu_n(G) \geq -\inf\{\Lambda^*(x); x \in G \cap \mathcal{F}\}$$

where \mathcal{F} is the set of exposed points of Λ^* whose exposed hyperplane lies in $\text{int } D_\Lambda$.

c) If Λ is essentially smooth, l.s.c., then we have a LDP of good rate function Λ^* .

DEFINITION 2.9. The sequence $\{Y_n\}$ is said to be exponentially tight if for each $\alpha > 0$ there exists a compact set K_α such that

$$\limsup \frac{1}{a_n} \log P_n(Y_n \in (K_\alpha)^c) \leq -\alpha.$$

2.4. VARADHAN THEOREM

We now study integrals of functions.

Let (Y_ϵ) be a family of r.v. with values in a regular topological space X , and (μ_ϵ) the corresponding distributions. We consider the following properties :

(LDP) (μ_ϵ) satisfies a LDP and its rate function I is good.

(LIM) For all $f \in \mathcal{C}_b$ the following limit exists :

$$\Lambda_f = \lim_{\epsilon} \epsilon \log \int e^{f(x)/\epsilon} d\mu_\epsilon(x).$$

THEOREM 2.10.

a) (LDP) \Rightarrow (LIM) with

$$(*) \quad \Lambda_f = \sup f - I.$$

b) (LIM) + exponential tightness implies (LDP) with

$$(**) \quad I(x) = \sup\{f(x) - \Lambda_f ; f \in \mathcal{C}_b\}$$

and (*) holds.

Part a) is Varadhan's theorem and b) is Bryc's theorem

We can weaken the conditions. Assume that (μ_ϵ) satisfies a LDP and that f is continuous. Assume moreover that the following uniform exponential integrability holds :

$$\lim_{M \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \epsilon \log E[e^{f(Y_\epsilon)/\epsilon} 1_{\{f(Y_\epsilon) \geq M\}}] = -\infty$$

then LIM is satisfied.

For instance, if

$$\limsup_{\epsilon \rightarrow 0} \epsilon \log E[e^{\gamma f(Y_\epsilon)}] < \infty$$

for some $\gamma > 1$, then we have the uniform exponential integrability.

2.5. SCHILDER'S THEOREM

Let B_t be a standard brownian motion in \mathbf{R} , $B_\epsilon(t) := \epsilon B_t$ and let (μ_ϵ) the family of corresponding distributions in $\mathcal{C}_0([0,1])$. Let $H_1 \subset \mathcal{C}_0([0,1])$ be the Cameron Martin space of absolutely continuous functions φ whose derivative $\dot{\varphi}$ is square-integrable.

THEOREM 2.11. *The family (μ_ϵ) satisfies a LDP of speed ϵ^{-2} and good rate function I defined on $\mathcal{C}_0([0,1])$ by :*

$$I(\varphi) = \int_0^1 [\dot{\varphi}(t)]^2 dt, \quad \varphi \in H_1$$

$$I(\varphi) = +\infty$$

in other cases .

There are various proofs of this result. The classical proof (Varadhan) uses for the lower bound a change of probability (Girsanov), and for the upper bound a discretization by polygonal lines . We can also proof this result as an application of a Cramer's theorem in infinite dimension.

This can be extended to diffusion processes with small parameter

$$dx_t^\epsilon = b(x_t^\epsilon)dt + \sqrt{\epsilon} \sigma(x_t^\epsilon) dw_t, \quad \sigma > 0.$$

This is the Freidlin-Wentzell theory (see Dembo-Zeitouni p.187-212). The action functional is then

$$I(\varphi) = \int_0^1 \frac{[\dot{\varphi}(t) - b(\varphi(t))]^2}{\sigma^2(\varphi(t))} dt.$$

3. BRANCHING PROCESSES

3.1. TREES

We give here some notations and background on branching processes (see Neveu (1986), Chauvin (1986), Lyons (1998)). If $(p_k)_{k \in \mathbf{N}}$ is a probability distribution, the process Z_n is defined by $Z_0 = 1$ (the ancestor) and this ancestor give birth to k descendents with probability p_k . These descendents have offspring independently with the same probability distribution.

Formally, let \mathbb{U} be the space of finite sequences $u = j_1 \dots j_n$ of strictly positives integers. The sequence \emptyset belongs to \mathbb{U} . So

$$\mathbb{U} = \{\emptyset\} \cup \bigcup_{n \geq 1} \mathbf{N}^{*n}$$

The length of $u \in \mathbb{U}$ is denoted by $|u|$; the concatenation of u and v is denoted by uv .

A tree ω is a subset of \mathbb{U} satisfying the following conditions :

- (i) $\emptyset \in \omega$
- (ii) $uv \in \omega \Rightarrow u \in \omega$
- (iii) $u \in \omega \Rightarrow \exists N_u(\omega) \in \mathbb{N}$ such that $uj \in \omega, j \geq 1 \Leftrightarrow 1 \leq j \leq N_u(\omega)$

Let N instead of N_\emptyset . The set Ω of trees is provided with the σ -field \mathcal{F} generated by $\Omega_u, u \in \mathbb{U}$ where $\Omega_u = \{\omega \in \Omega, u \in \omega\}$. So N_u are r.v. defined on Ω_u . For every integer n the n -th generation is

$$z_n(\omega) = \{u \in \omega, |u| = n\}$$

and

$$Z_n(\omega) = \text{Card } z_n(\omega)$$

(with $Z_1 = N$).

The set of trees Ω is now provided with a filtration \mathcal{F}_n generated by $z_m, 0 \leq m \leq n$ and with (measurable) shifts

$$\begin{aligned} T_u : \Omega &\rightarrow \Omega \\ \omega &\mapsto T_u(\omega) = \{v, v \in U, uv \in \omega\} \end{aligned}$$

so that for all n et k ,

$$(1.1) \quad Z_{n+k} = \sum_{u \in z_n} Z_k \circ T_u$$

We now make (Ω, \mathcal{F}) a probability space.

THEOREM 3.1. *Let $p = (p_k)_{k \geq 0}$ a probability distribution. There exists a unique probability P on (Ω, \mathcal{F}) such that*

- (i) *the distribution of N is p .*
- (ii) *conditionally on $\{N = j\}$, the r.v. T_1, T_2, \dots, T_j are independent and P -distributed.*

Consequently, for all $n \geq 0$, conditionally on \mathcal{F}_n , the r.v. $\{T_u, u \in z_n\}$ are independent and P -distributed. This means that for all measurable positive functions $f_u, u \in U$

$$E^{\mathcal{F}_n} \left(\prod_{u \in z_n} f_u \circ T_u \right) = \prod_{u \in z_n} E(f_u) \quad (3.1)$$

The sequence $(Z_n)_{n \geq 0}$ is called a Galton-Watson process of offspring distribution p and initial state $Z_0 = 1$. The formula (3.1) is called the branching property.

The classical tool is the generating function $f(s) = \sum s^k p_k$.

3.2. MARKED TREES

The idea is to mark each node of a tree with a mark.

Branching random walk

We mark a node u with a point process $(X_{u1}, \dots, X_{u\nu_u})$ representing the positions of the children of u relatively to the position of u .

Branching diffusion

We mark a node u with a trajectory.

Multitype processes

We mark a node u with an element of a finite set of types $E = \{1, \dots, r\}$.

We can of course mix the models to obtain the multitype branching random walks or multitype branching diffusions. In these lecture, marks are supposed to fit with the branching property and markovian along branches (see Rouault 1987).

In each model, at the end of the construction each node u of a random tree is provided with a label Y_u in some set S . Important tools are then the generating functional M acting on φ , function of S in $[0, 1]$ by :

$$M(\varphi)(x) = E_x \prod_{u \in z_1} \varphi(Y_u). \quad (3.2)$$

and the additive functional

$$S(\varphi)(x) = E_x \sum_{u \in z_1} \varphi(Y_u). \quad (3.3)$$

4. LARGE DEVIATIONS AND GALTON-WATSON PROCESSES

We present here two types of applications of large deviations in supercritical Galton-Watson process processes. The first one is a large deviation theorem for the Lotka-Nagaev estimator of the mean. The second one is particularly nice. It uses the "self-similarity" of the r.v. W limit of the process to give tail behaviours of the distribution of W (in 0 and in ∞). Both need some properties of the iterated generating functions.

4.1. ASYMPTOTIC BEHAVIOUR OF THE GENERATING FUNCTION

When $p_0 = 0$ and $p_1 \neq 0$ then

$$p_1^{-n} f_n(s) \rightarrow Q(s) \quad (4.1)$$

for $0 \leq s < 1$.

When there is a finite maximum family size d , it is known that

$$\frac{\log f_n(s)}{d^n} \rightarrow \log B_r(s) \quad (4.2)$$

for $s > 1$ where B_r is the right Böttcher function (Kuczma Choczewski and Ger 1990).

When there is a minimum $k \geq 2$ family size, it is known that

$$\frac{\log f_n(s)}{k^n} \rightarrow \log B_l(s) \quad (4.3)$$

for $0 \leq s < 1$ where B_r is the left Böttcher function.

The three limit functions, Q , $\log B_r$ and $\log B_l$ satisfy a Schröder equation :

$$F \circ f = \lambda F \quad (4.4)$$

with $\lambda = p_1$, d and k respectively.

4.2. GROWTH RATES

In a supercritical Galton-Watson process with $p_0 = 0$ the classical Lotka-Nagaev estimator of the mean m is $\frac{Z_{n+1}}{Z_n}$ (Dion 1991). We present here large deviations of this estimator essentially due to Athreya.

THEOREM 4.1.

Let $p_0 = 0, p_1 > 0, p_j \neq 1$ for $j \geq 1$ and $m > 1$. Let $E(Z_1^{2r+\delta}) < \infty$ for some $r \geq 1$ and $\delta > 0$ such that $p_1 m^r > 1$. Then

$$\lim_{n \rightarrow \infty} p_1^{-n} P(|\frac{Z_{n+1}}{Z_n} - m| > \epsilon) = g(\epsilon) < \infty$$

This last result leads to a LDP for $\frac{Z_{n+1}}{Z_n} - m$ at rate n^{-1} with rate function

$$I(x) = -\log p_1 \text{ if } x \neq 0, \quad I(0) = 0$$

This is indeed a rate function, but it is not *good*.

THEOREM 4.2.

In the minimum family size case, let $\Lambda(\theta) = \log f(e^\theta)$. If there is some $\theta > 0$ in dom Λ , then the family $\frac{Z_{n+1}}{Z_n}$ satisfies a LDP at speed k^n with good rate function $\log B_l(e^{-\Lambda^*(\cdot)})$

Proof. : The upper bound is proved by Athreya (th. 3). We give it for the sake of completeness. Let F be a closed set. Conditioning on Z_n gives

$$P(\frac{Z_{n+1}}{Z_n} \in F) = \sum_j P(Z_n = j) P(\frac{S_j}{j} \in F) \quad (4.5)$$

where S_k is the sum of k independent r.v. distributed as Z_1 . The Chernov upper bound (2.3) yields

$$P(|\frac{Z_{n+1}}{Z_n} \in F) \leq 2f_n(\exp -\Lambda^*(F))$$

and the upper bound follows. For the lower bound, let us first remark that in (4.5) the sum actually begins at $j = k^n$. Let $\alpha > 0$. From the Chernov lower bound (2.2), there exists N such that for $j > N$

$$P\left(\frac{S_j}{j} \in F\right) \geq \exp -[\Lambda^*(G) + \alpha]$$

This yields for $k^n > N$,

$$P\left(\frac{Z_{n+1}}{Z_n} \in G\right) \geq f_n(\exp -\Lambda^*(G) - \alpha)$$

and then

$$\liminf k^{-n} \log P\left(\frac{Z_{n+1}}{Z_n} \in G\right) \geq \log B_l \circ (\exp -\Lambda^*(G) - \alpha)$$

Since it holds for all α the lower bound holds.

These results may be extended to the multitype case (see Athreya and Vidyashankar 1995).

4.3. TAIL BEHAVIOUR

We study here the tail behaviour of the r.v. $W := \lim_n \frac{Z_n}{m^n}$ associated to a supercritical Galton-Watson process. Let Φ the Laplace transform of W defined by $\Phi(s) = Ee^{sW}$. It satisfies the equation :

$$f \circ \Phi(s) = \Phi(ms). \quad (4.6)$$

THEOREM 4.3.

1) In the finite maximum size case (f is a polynomial of degree d), let

$$\gamma = \frac{\log d}{\log m} (> 1) \quad \text{and} \quad \frac{1}{\gamma} + \frac{1}{\gamma^*} = 1.$$

a) As $s \rightarrow \infty$

$$\log \Phi(s) = \log B_r \circ \Phi(s) + O(1) \quad (4.7)$$

and the function H defined by $H(s) = s^{-\gamma} \log B_r \circ \Phi(s)$, is multiplicatively periodic of period m , positive and continuous.

b) As $x \rightarrow \infty$

$$-\log P(W \geq x) = (\log B_r \circ \Phi)^*(x) + o(x^{\gamma^*})$$

and $(\log B_r \circ \Phi)^*(x) = x^{\gamma^*} H^\dagger(x)$ where H^\dagger is multiplicatively periodic with period $m^{\gamma-1}$.

c) As $x \rightarrow \infty$

$$\liminf \frac{-\log P(W \geq x)}{x^{\gamma^*}} = \frac{1}{\gamma^*(\gamma\tau)^{\frac{1}{\gamma-1}}}$$

$$\limsup \frac{-\log P(W \geq x)}{x^{\gamma^*}} = \frac{1}{\gamma^*(\gamma\underline{\tau})^{\frac{1}{\gamma-1}}}$$

where $\underline{\tau} = \min_{x \leq m} H(x)$ and $\overline{\tau} = \max_{x \leq m} H(x)$

2) In the minimum size case, let

$$\delta = \frac{\log k}{\log m} (< 1) \quad \text{and} \quad \frac{1}{\delta} - \frac{1}{\delta^*} = 1.$$

As $x \rightarrow 0$

$$-\log P(W \leq x) = (\log B_l \circ \Phi)^*(x) + o(x^{-\delta^*})$$

and $(\log B_l \circ \Phi)^*(x) = x^{\delta^*} L^\dagger(x)$ where L^\dagger is multiplicatively periodic with period $m^{1-\delta}$.

1) a) is due to Harris (1948).

1) b) and 2) are from Biggins and Bingham (1993). These results can be proved by application of the Gärtner-Ellis theorem.

Actually from (4.6) we have

$$\Phi(m^n s) = f_n \circ \Phi(s) \tag{4.8}$$

and from (4.2) we deduce that $\Lambda_n(\theta) := \frac{1}{m^{n\gamma}} \log E \exp \theta m^n W$ if $\theta > 0$, has the limit $\Lambda(\theta) = \log B_r \circ \Phi(\theta)$. So we can apply a one-sided version of the Gärtner-Ellis theorem yielding a one-sided LDP for the family of distributions of $(m^{n(1-\gamma)} W)$ at speed $m^{-n\gamma}$ with rate function $(\log B_r \circ \Phi)^*$.

c) is a consequence of b) and results of Liu (see p.63).

To prove 2), let L defined for $s \geq 0$ by :

$$L(s) = -s^{-\delta} \log B_l \circ \Phi(-s).$$

It is straightforward from (4.4) and (4.6) to see that L is multiplicatively m -periodic, continuous and positive, and that

$$\frac{\log \Phi(-m^n s)}{k^n} \rightarrow -s^\delta L(s).$$

From this we deduce that $\Lambda_n(\theta) := \frac{1}{m^{n\delta}} \log E \exp \theta m^n W$ if $\theta < 0$, has the limit $\Lambda(\theta) = \log B_l \circ \Phi(\theta)$. Again a one-sided version of the Gärtner-Ellis theorem yields a LDP for the family of distributions of $(m^{n(1-\delta)} W)$ at speed $m^{-n\delta}$ with rate function $(\log B_l \circ \Phi)^*$.

REMARK. For the tree martingale issued from more general branching processes there are also tail results (see Liu (1996)). For instance, for the cascade process (see Waymire (1998)) the rate of increasing of $-\log P_r(Z_\infty > x)$ is $O(x^{\gamma^*})$ with $\gamma^* = \frac{\log b \|W\|_\infty}{\log N}$.

5. LARGE DEVIATIONS IN SPATIAL PROCESSES

As shortly presented in the introduction the competition between small probabilities and large populations provides with three problems :

the speeds of propagation (borders separating supercritical and subcritical ranges)

the growth rates (of population) in the supercritical range

the evaluation of presence probabilities in the subcritical range.

5.1. RATE OF PROPAGATION

A) Branching Random Walk

Marks of nodes are point processes. We suppose these points processes i.i.d. of generating functional ψ so that

$$M(\varphi)(x) = \psi(\varphi(\cdot - x)), \quad (5.1)$$

(see (3.2)) and of intensity measure μ so that

$$S(\varphi(x)) = \int \varphi(y - x) d\mu(x) \quad (5.2)$$

(see (3.3)). The offspring mean is $m = \int d\mu(x)$. We assume $m > 1$ (the underlying Galton-Watson is supercritical). Let Λ be the log-Laplace of μ . We assume that $0 \in \text{intdom } \Lambda$. We assume also for the sake of simplicity that μ is zero-mean.

For $n > 0$ we denote \mathcal{F}_n the sigma field generated by the marks of nodes of length smaller than n . Let for $u \in \omega$, $S_u = X_{u_1} + X_{u_1 u_2} + \dots + X_{u_1 u_2 \dots u_n}$. If δ_a is the Dirac mass in $a \in \mathbb{R}$, let $Z_0 = \delta_0$ and for $n > 0$

$$Z_n = \sum_{u \in z_n} \delta_{S_u} \quad ,$$

so that if B is a Borel set of \mathbb{R} , we have:

$$Z_n(B) = \text{card } \{u \in z_n : S_u \in B\}.$$

For all $\theta \in \mathbb{R}$ and $n > 0$, we denote

$$W_n(\theta) = \sum_{u \in z_n} \exp[\theta S_u - n\Lambda(\theta)].$$

This is a \mathcal{F}_n - martingale positive. Let $W(\theta)$ be its a.s. limit. From our assumptions, it turns out that $\{x : \Lambda^*(x) \leq 0\}$ is a compact set $[c^-, c^+]$.

THEOREM 5.1.

If

$$M_n^+ = \max\{S_u; u \in z_n\} \quad , \quad M_n^- = \min\{S_u; u \in z_n\}$$

then a.s. as $n \rightarrow +\infty$: $\lim M_n^+/n = c^+$ and $\lim M_n^-/n = c^-$.

Proof.

We will only treat the case of the maximum.

a) Let us show first that a.s. $\limsup M_n/n \leq c^+$.

For all a we have : $\{Z_n([na, +\infty[) \geq 1\} = \{M_n \geq na\}$ and then $P\{M_n \geq na\} \leq EZ_n([na, +\infty[)$ by Markov inequality. For all f measurable positive,

$$\begin{aligned} Z_n(f) &= \int f(x) dZ_n(x) = \sum_{u \in z_n} f(S_u) \\ &= \sum_{u \in z_{n-1}} \sum_{v \in z_1^u} f(S_u + X_{uv}). \end{aligned}$$

Hence

$$E[Z_n(f) | \mathcal{F}_{n-1}] = Z_{n-1} \star \mu(f).$$

This yields :

$$E(Z_n(g)) = \mu^{\star n}(g).$$

In particular $EZ_n([na, +\infty[) = \mu^{\star n}([na, +\infty[)$ and from (2.3)

$$\mu^{\star n}([na, +\infty[) \leq \exp -n\Lambda^*(a) \text{ if } a > 0.$$

Moreover if $\Lambda^*(a) > 0$, then $\sum P(Z_n([na, +\infty[) \geq 1) < \infty$ and by Borel-Cantelli there exists a.s. n_0 such that $Z_n([na, +\infty[) = 0$ for all $n \geq n_0$. This gives a.s. $\limsup M_n/n \leq c^+$.

b) The proof of the lower bound is an easy consequence of the next result. □

PROPOSITION 5.2.

If $a > 0$ et $\Lambda^*(a) < 0$ then a.s.

$$\lim \frac{1}{n} \log Z_n([na, +\infty[) = -\Lambda^*(a).$$

Proof. The upper bound is proved using Borel-Cantelli and argument from a). The lower bound can be proved using an auxiliary branching process (Biggins 1976) or using the following martingale argument (Chauvin, 1986, see also Neveu 1988).

LEMMA 5.3.

1) For $\theta \in \mathbb{R}$, $W_n(\theta)$, $n > 0$ is a \mathcal{F}_n - martingale positive. Let $W(\theta)$ be its a.s. limit.

2) For all a such that $\Lambda^*(a) < 0$, let $\theta = \Lambda'(a)$. Then $W_n(\theta) \rightarrow W(\theta)$ in \mathbb{L}^1 and $P(W(\theta) = 0) = 0$.

3) If $\Lambda^*(a) > 0$, then $W_n(\theta) \rightarrow 0$ a.s. when $n \rightarrow \infty$.

Suppose this lemma proved. We have

$$W_n(\theta) = \Gamma_n + \sum_{u \in z_n: S_u \in [n(a-\epsilon), n(a+\epsilon)]} \exp[\theta S_u - n\Lambda(\theta)].$$

In the last sum we can bound above S_u by $n(a + \epsilon)$ ($\theta > 0$) and keep only those u such that $S_u \in [n(a - \epsilon), +\infty[$. This yields

$$W_n(\theta) \leq Z_n([n(a - \epsilon), +\infty[) \exp[n(a + \epsilon)\theta - n\Lambda(\theta)] + \Gamma_n.$$

If

$$d\mu_\theta(x) = \exp \theta x - \Lambda(\theta) d\mu(x)$$

then

$$\begin{aligned} E\Gamma_n &= \int_{x \notin [n(a - \epsilon), n(a + \epsilon)]} \exp[\theta x - n\Lambda(\theta)] d\mu^{*n}(x) \\ &= 1 - \mu_\theta([n(a - \epsilon), n(a + \epsilon)]) \end{aligned}$$

From (2.1) this series converges and then a.s. $\Gamma_n \rightarrow 0$. From the lemma,

$$\underline{\lim}_n Z_n([n(a - \epsilon), +\infty[) \exp[n\theta(a + \epsilon) - n\Lambda(\theta)] \geq W(\theta) > 0$$

a.s.. For all $a > 0$ such that $\Lambda^*(a) < 0$ and all $\epsilon > 0$, we then have :

$$\underline{\lim}_n \frac{1}{n} \log Z_n([n(a - \epsilon), +\infty[) \geq -\Lambda^*(a) - \epsilon\theta.$$

It means that for any $a > 0$ such that $\Lambda^*(a) < 0$ and any ϵ small enough,

$$\underline{\lim}_n \frac{1}{n} \log Z_n([na, +\infty[) \geq -\Lambda^*(a + \epsilon) - \epsilon\Lambda'(a + \epsilon).$$

Then let $\epsilon \rightarrow 0$.

Proof of 5.3. :

See Lyons (1997). □

DEFINITION 5.4. We assume $0 \in \text{int dom } \Lambda$. We say that $a \in \mathbb{R}$ is in the range of supercriticality if $a \in \Lambda'(\text{int dom } \Lambda)$ and if $\Lambda^*(a) < 0$. Let θ such that $\Lambda'(\theta) = a$. We say that a is in the range of subcriticality if $a \in \Lambda'(\text{int dom } \Lambda)$ and if $\Lambda^*(a) > 0$.

B) Malthusian phenomena

A classical model for growing populations is the Bellman-Harris process. Each individual has a (random) lifetime of distribution G and at its death time give birth to a random number (of mean m) of children. The population alive at time t comes from different generations. For large t , the empirical mean of lifetimes on a given branch of the genealogic tree is approximately $\int s dG(s)$ from the LLN. Branches corresponding to means less than $\int s dG(s)$ are less probable, but had time to give more offsprings. There is a compensation and asymptotically the number of alive individual is near $e^{\alpha t} W$. The Malthusian parameter α is the unique positive solution of

$$\int_0^\infty e^{-\alpha t} m dG(t) = 1$$

In general this very classical result (Athreya-Ney 1971 p.) is proved using the renewal theory. Here we give the "large deviations heuristics" to allow a generalization.

Let X_1, \dots, X_n, \dots independent and G -distributed r.v.. For every $\beta > 0$, the contribution of generation number $[t/\beta]$ to the mean population alive at time t is roughly (for t large)

$$\begin{aligned} m^{[t/\beta]} P(X_1 + \dots + X_{[t/\beta]} \simeq t) &= m^{[t/\beta]} \exp\left(-\frac{t}{\beta} \Lambda_G^*(\beta) + o(1)\right) \\ &= \exp\left(-\frac{t}{\beta} (\Lambda_G^*(\beta) - \log m + o(1))\right). \end{aligned}$$

When β takes different values, we get exponentials of different orders. The larger one is overwhelming. It is precisely $\exp \alpha' t$ where

$$\alpha' = - \inf_{\beta > 0} \left(-\frac{1}{\beta} (\Lambda_G^*(\beta) - \log m)\right)$$

This infimum is reached in $\beta_0 = \int_0^\infty s e^{-\alpha s} m dG(s)$ and $\alpha' = \alpha$. The generations whose number is $\frac{t}{\beta_0} + o(t)$ are overwhelming (look at the means and apply a LLN). This β_0 is the mean age at childbearing, well known in demography. (see Jagers p.214).

This heuristics allows us to extend it to the so called Crump-Mode-Jagers branching process with random walk. We mark every individual y with a lifelength λ_y , a birthtime T_y and a (fixed) position X_y . Its immediate offspring is described by a point process ξ_y on $\mathbb{R}^+ \times \mathbb{R}$. Each point yk of ξ_y corresponds to a child of y , the first coordinate is the age of y when yk was born and the second coordinate is the increment of position from y to yk . Here we assume that given the tree, all the (λ_y, ξ_y) are i.i.d.. The jumps along a genealogical branch are $\mathbb{R}^+ \times \mathbb{R}$ valued. The asymptotic scaling of Bellman-Harris' model ($t \rightarrow \infty$) leads us to introduce a small parameter ϵ and perform a scaling of rate ϵ in $\mathbb{R}^+ \times \mathbb{R}$. In other words, from a reference model, we build a family of ϵ indexed processes, multiplying the jumps (in $\mathbb{R}^+ \times \mathbb{R}$) and the lifetimes by ϵ . Let $\zeta_t^\epsilon(A)$ the size of the population alive at t and located in A , Borel set of \mathbb{R} . Let μ be the intensity of the point process, Λ its log-Laplace, assumed to be finite everywhere and Λ^* its Cramer transform.

Let α 1-homogeneous from $\mathbb{R}^+ \times \mathbb{R}$ to \mathbb{R}^+ , by

$$\alpha(t, x) = - \inf_{\beta > 0} \frac{1}{\beta} \Lambda^*(t\beta, x\beta).$$

We showed (Laredo-Rouault 1983), under some assumptions, that there exists a cone \mathcal{S} such that for every interval I of \mathbb{R}

- 1) If $t \times I \in \mathbb{R} - \hat{\mathcal{S}}$ then a.s. $\zeta_t^\epsilon(I) = 0$ for ϵ small enough.
- 2) If $t \times I \cap \mathcal{S}$ is non empty, then as $\epsilon \rightarrow 0$,

$$\epsilon \log \zeta_t^\epsilon(I) \rightarrow \sup_{x \in I} \alpha(x, t)$$

in probability.

Function α plays the role of a local Malthusian parameter, and relation ? becomes

$$\Lambda\left(-\frac{\partial\alpha}{\partial t}, -\frac{\partial\alpha}{\partial x}\right) = 0.$$

5.2. RANGE OF SUPERCRITICALITY

We improve here the result of theorem 5.1 , giving sharp estimates of the population near na where a is in the range of supercriticality.

Single type case

From the classical result of Bahadur-Rao (see Dembo-Zeitouni p.95), we get :

$$EZ_n(na + I_\delta) = \mu^{*n}(na + I_\delta) = \frac{\exp -n\Lambda^*(a) e^{\theta\delta} - e^{-\theta\delta}}{\sqrt{2\pi\Lambda''(\theta)} \theta} (1 + o(1)).$$

where $I_\delta = [-\delta, +\delta]$. The following result is due to Biggins (1979).

THEOREM 5.5. *If a is in the range of supercriticality and if the moment condition*

$$E[W_1(\theta) \log_+^\epsilon W_1(\theta)] < \infty \text{ for some } \epsilon > 5/2$$

holds, then a.s.

$$\frac{Z_n(na + I_\delta)}{EZ_n(na + I_\delta)} \rightarrow W(\theta)$$

Multitype case

The result is due to Bramson, Ney and Tao (1992). It is stated in the case of a lattice random walk.

Let $Z_{ij}^n([x])$ be the number of type j particles at the point "integer part of x " at time n , descended from a type i ancestor at time 0. Let $M = (m_{ij})$ the matrix of intensity measures of this process. We suppose it irreducible and aperiodic. For $\theta \in \mathbb{R}$, let

$$\hat{Z}_{ij}^n(\theta) = \sum_{x \in I} e^{\theta x} Z_{ij}^n(x)$$

and

$$\hat{M}(\theta) = (\hat{m}_{ij}(\theta)) = \sum_{x \in I} e^{\theta x} m_{ij}(x) ; i, j \in S$$

Let $\lambda(\theta)$ be the Perron-Frobenius root of $\hat{M}(\theta)$ and $\Lambda(\theta) = \log \lambda(\theta)$. Let also $u(\theta)$ and $v(\theta)$ the normalized right and left eigenvectors associated with $\lambda(\theta)$. It is easy to see that

$$W_i^n(\theta) = \frac{\sum_j \hat{Z}_{ij}^n(\theta) v_j(\theta)}{\lambda(\theta)^n v_i(\theta)} \quad (5.3)$$

is a positive martingale. Let $W_i(\theta)$ be its limit. We assume $\lambda(0) > 1$ which means that the underlying multitype Galton-Watson process is supercritical. We assume that a is in the range of supercriticality. Moreover we assume the moment condition :

$$E(\hat{Z}_{ij}(\theta))^p < \infty \text{ for some } p > 1 \text{ for } i, j = 1, \dots, d.$$

THEOREM 5.6. *Under the above assumptions, for $i, j = 1, \dots, d$ a.s.*

$$\lim \sqrt{2\pi n} \sigma_i(\theta) \exp\{\theta([an] - an)\} \exp n\Lambda^*(a) Z_{ij}^n([an]) = u_j(\theta)v_i(\theta)W_i(\theta)$$

where $\sigma_i(\theta)$ is an appropriate second moment parameter.

COROLLARY 5.7. *For $i, j, k = 1, \dots, d$, a.s.*

$$\frac{Z_{ij}^n([an])}{Z_{jk}^n([an])} \rightarrow \frac{u_j(\theta)v_i(\theta)}{u_k(\theta)} v_j(\theta).$$

5.3. RANGE OF SUBCRITICALITY

We want here estimates of probability of presence in the range of subcriticality. Let us first recall the classical result for the subcritical Galton-Watson process (Athreya Ney 1972). We have, as $n \rightarrow \infty$,

$$\frac{P(Z_n > 0)}{EZ_n} \rightarrow C$$

and

$$C \neq 0 \Leftrightarrow \sum (k \log k) p_k < \infty$$

Let us recall the main ideas of the proof to extend it further. If f is the g.f. of the offspring (with mean $m < 1$), then the sequence

$$u_n = P(Z_n > 0) = 1 - f_n(0)$$

satisfies

$$\begin{aligned} u_{n+1} &= 1 - f(1 - u_n) \\ u_0 &= 1 \end{aligned} \tag{5.4}$$

We compare it with the sequence

$$v_n = EZ_n = m^n$$

which satisfies obviously

$$\begin{aligned} v_{n+1} &= mv_n \\ v_0 &= 1 \end{aligned} \tag{5.5}$$

We introduce the classical auxiliary function :

$$r(s) = m - \frac{1 - f(1 - s)}{s}$$

which is increasing from 0 and we deduce :

$$u_n = v_n \prod_{j=0}^{n-1} \left[1 - \frac{r(u_j)}{m}\right] \quad (5.6)$$

and of course

$$u_n \leq v_n$$

It is known that for all $\lambda < 1$

$$\sum_j r(O(\lambda^j)) < \infty \iff \sum (k \log k) p_k < \infty$$

A) In the Branching Random Walk We may follow the same route . Let

$$u_n(x) = P_x(Z_n(I_\delta) \neq 0)$$

and

$$v_n(x) = E_x Z_n(I_\delta)$$

Let us remark that since I_δ is symmetric $u_n(x) = P_{-x}(Z_n(I_\delta) \neq 0)$ and since the motion is homogeneous $u_n(x) = P(Z_n(x + I_\delta) \neq 0)$, and $v_n(x) = E Z_n(x + I_\delta)$. With the branching property, we can prove :

$$\begin{aligned} u_{n+1} &= 1 - \Psi[1 - u_n] \\ u_0 &= 1_{I_\delta} \end{aligned} \quad (5.7)$$

and

$$\begin{aligned} v_{n+1} &= v_n \star \mu \\ v_0 &= 1_{I_\delta} \end{aligned} \quad (5.8)$$

(So-called M-equation and S-equation of Ikeda, Nagasawa and Watanabe).

THEOREM 5.8.

If a is in the range of subcriticality and if

$$EW_1(\theta) \log_+^{1+\epsilon} W_1(\theta) < \infty \text{ and } EW_1(0) \log_+^{1+\epsilon} W_1(0) < \infty$$

then

$$\frac{P(Z_n(nx + I_\delta) \neq 0)}{EZ_n(nx + I_\delta)} \rightarrow C > 0.$$

This result is in Rouault (1993). The proof is built on the following representation formulas :

$$\begin{aligned} u_n(x) &= m^n E_x \{u_0(S_n) \\ &\prod_{k=1}^n [1 - \int_{\mathcal{M}} \int_0^1 [1 - \exp \int_{\mathbb{R}} \log 1 - \lambda u_{n-k}(S_{k-1} + y) Z(dy)] d\lambda P^{1X_k}(dZ)]\} \\ v_n(x) &= m^n E_x u_0(S_n) \end{aligned} \quad (5.9)$$

where $\{S_n\}$ is a random walk $m^{-1}\mu$ -distributed.

B) In the homogeneous branching brownian motion

Here recursion formulas become p.d.e.. Let

$$u(t, x) := P_x(Z_t(]-\infty, 0]) > 0) = P_0(Z_t([x, +\infty[> 0)$$

(by symmetry) and

$$v(t, x) = E_x Z_t(]-\infty, 0]) = E_0 Z_t([x, +\infty[)$$

Using a continuous version of the branching property, we get the M-equation, which is the famous Kolmogorov-Petrovski-Piscounov equation (KPP) :

$$\begin{aligned} \frac{\partial u}{\partial t} &= \frac{1}{2} \frac{\partial^2 u}{\partial x^2} + \alpha[1 - u - f(1 - u)] \\ u(0, \cdot) &= 1_{]-\infty, 0]}. \end{aligned} \quad (5.10)$$

The S-equation is :

$$\begin{aligned} \frac{\partial v}{\partial t} &= \frac{1}{2} \frac{\partial^2 v}{\partial x^2} + \alpha(m - 1)v \\ v(0, \cdot) &= 1_{]-\infty, 0]}. \end{aligned} \quad (5.11)$$

We can use the representation with an auxiliary brownian motion :

$$v(t, x) = E 1_{]-\infty, 0]}(x + W_t) \exp \alpha(m - 1)t$$

and by the Feynmann-Kac formula :

$$u(t, x) = E \left[1_{]-\infty, 0]}(x + W_t) \exp \alpha(m - 1)t - \int_0^t \alpha r(u(t - s, W_s)) ds \right]$$

THEOREM 5.9. For $c > c_0 = \sqrt{2\alpha(m - 1)}$ and under the $(k \log k)$ assumption, we have

$$\lim_{\substack{x \rightarrow \infty, t \rightarrow \infty \\ x/t \rightarrow c}} \frac{u(t, x)}{v(t, x)} > 0$$

and

$$v(t, x) \sim \frac{e^{-\frac{t}{2}(c^2 - c_0^2)}}{c\sqrt{2\pi t}}$$

The rates of propagation are $\pm c_0 = \pm \sqrt{2\alpha(m - 1)}$ and $|c| > c_0$ defines the range of subcriticality.

C) In the inhomogeneous BBM

The model consists in a brownian motion with a small variance (ϵ^2), a binary splitting (to simplify) and a branching rate $\frac{c(\cdot)}{\epsilon^2}$ with $c \in \mathcal{C}^3(\mathbb{R}, \mathbb{R}^+)$.

We study $u^\epsilon(t, x) = P_x^\epsilon\{Z_t([\!-\infty, 0]) \geq 1\}$ and $v^\epsilon(t, x) = E_x^\epsilon\{Z_t([\!-\infty, 0])\}$.
The M-equation is :

$$\begin{aligned} u_t^\epsilon &= \frac{\epsilon^2}{2} u_{xx}^\epsilon + \frac{c(x)}{\epsilon^2} u^\epsilon (1 - u^\epsilon) \\ u^\epsilon(0, \cdot) &= 1_{\mathbb{R}^-} \end{aligned} \quad (5.12)$$

and the S-equation

$$v_t^\epsilon = \frac{\epsilon^2}{2} u_{xx}^\epsilon + \frac{c(x)}{\epsilon^2} u^\epsilon. \quad (5.13)$$

We use the representations :

$$\begin{aligned} u^\epsilon(t, x) &= E[g(x + \epsilon B_t) \exp \frac{1}{\epsilon^2} \int_0^t c(x + \epsilon B_s) [1 - u^\epsilon(t - s, x + \epsilon B_s)] ds] \\ v^\epsilon(t, x) &= E[g(x + \epsilon B_t) \exp \frac{1}{\epsilon^2} \int_0^t c(x + \epsilon B_s) ds]. \end{aligned}$$

THEOREM 5.10.

$$\epsilon^2 \log u^\epsilon(t, x) \rightarrow V^*(t, x)$$

where

$$V^*(t, x) = \inf_{\tau} \sup \left\{ \int_0^{t \wedge \tau} [c(\phi_s) - \frac{1}{2} \dot{\phi}_s^2] ds; \phi_0 = x, \phi_t \in \text{Supp } g \right\}$$

and the supremum is over all stopping times.

The range of subcriticality is $\{(t, x) : V^*(t, x) < 0\}$. There exists also another formula for V^* :

$$V^*(t, x) = \sup \left\{ \inf_{0 \leq a \leq t} \int_0^a [c(\phi_s) - \frac{1}{2} \dot{\phi}_s^2] ds; \phi_0 = x, \phi_t \in \text{Supp } g \right\}$$

We present now a sharp (i.e. non logarithmic) asymptotic formula. We assume c at most linear at infinity, and

(H1) The sup in V is reached in a unique path ϕ .

(H2) ϕ is a nondegenerate maximum.

That means that the operator A on $H^0 = \{\phi \in H^1 : \phi_0 = 0\}$ given by

$$\langle Ah, h \rangle = \int_0^t c''(\phi_s) h_s^2 ds$$

has its spectrum strictly bounded above by 1. We may define

$$C(\phi) = E \exp \frac{1}{2} \int_0^t c''(\phi_s) B_s^2 ds = \det(I - A)^{-\frac{1}{2}}.$$

For the bridge we define $C_0(\phi)$.

THEOREM 5.11.

Under H1 and H2,

a) If $\phi_t = 0$ and $p =: -\dot{\phi}_t > 0$

$$v^\epsilon(t, x) = \epsilon \exp \frac{V(t, x)}{\epsilon^2} p^{-1} (2\pi t)^{-1/2} C_0(\phi) (1 + o(1))$$

If $\phi_t < 0$

$$v^\epsilon(t, x) = \exp \frac{V(t, x)}{\epsilon^2} C(\phi) (1 + o(1)).$$

Let us give other assumptions :

(H3) The set $\{(\phi, a) : \phi_t \leq 0, \phi_0 = x \text{ and}$

$$\int_0^a [c(\phi_s) - \frac{1}{2} \dot{\phi}_s^2] ds = \inf_{0 \leq b \leq t} \int_0^b [c(\phi_s) - \frac{1}{2} \dot{\phi}_s^2] ds = V^*(t, x)\}$$

is a singleton.

This assumption implies in particular :

a) $V(t, x) = V^*(t, x)$

b) There exists one unique ϕ reaching the sup in V^*

c) The optimal path stays always ahead of the nonlinear front.

Under H1+H2+H3 we have

$$\phi_t = 0, \quad p > 0 \text{ et } p \geq [2c(0)]^{1/2} \text{ et } V = V^*.$$

We need

(H4) $p > [2c(0)]^{1/2}$.

THEOREM 5.12. If $V^*(t, x) < 0$, under assumptions H1, H2, H3, H4, we have :

$$u^\epsilon(t, x) = \epsilon \exp \frac{V(t, x)}{\epsilon^2} p^{-1} (2\pi t)^{-1/2} C_0(\phi) C_1(p) (1 + o(1))$$

where $C_1(p)$ depends only on p .

There are two extensions of this result. If the brownian motion is replaced by a non-degenerate diffusion (without drift to simplify) (see Rouques 1997), the functional is obtained by changing $\int \frac{1}{2} \dot{\phi}_s^2 ds$ in $\int \frac{1}{2} \frac{\dot{\phi}_s^2}{\sigma^2(\phi_s)} ds$. The brownian \mathbb{R}^d -case with a convenient initial condition is studied in Cohen-Rossignol 1997. In both cases the authors get new constants.

ACKNOWLEDGEMENTS

Many thanks to Prof. N. Yanev and the Bulgarian Academy of Sciences for the invitation to the Sozopol School, and for the warm hospitality. Many thanks to all the participants for stimulating discussions.

REFERENCES

- ATHREYA K.B. (1994) , Large deviation rates for branching processes-I. Single type case, *Annals of Appl. Prob.* **4** 779-790.
- ATHREYA, K.B. and NEY P. (1972) , Branching Processes, Springer.
- ATHREYA K.B. and VIDYASHANKAR, A. (1995) , Large deviation rates for branching processes-II. Multitype case, *Annals of Appl. Prob.* **5** 566-576.
- ATHREYA K.B. and VIDYASHANKAR, A. (1993) , Large Deviation Results for Branching Processes. *A Festschrift in Honour of Gopinath Lallianpur*, Springer.
- ATHREYA K.B. and VIDYASHANKAR, A. (1996) , Large deviation rates for supercritical and critical branching processes "Classical and Modern Branching Processes", *IMA 84, Athreya and Jagers, Eds* Springer.
- BEN AROUS G. and ROUAULT A. (1993) Laplace asymptotics for reaction diffusion equations *Probab. Theory and Rel. Fields* , **97** 259-285.
- BIGGINS, J.D. (1977) Chernoff's theorem in the branching random walk. *J. Appl. Probab.* **14** 630-636.
- BIGGINS, J.D. (1977) Martingale convergence in the branching random walk *J. Appl. Probab.* **14** 25-37.
- BIGGINS, J.D. (1978) The asymptotic shape of the branching random walk *Adv. Appl. Probab.* **10** 62-84.
- BIGGINS, J.D. (1979) Growth rates in the branching random walk *Z. Wahrschein. verw. Geb.* **48** 17-34.
- BIGGINS, J.D. (1980) Spatial spread in branching processes *Lect. Notes Biomath.* **38** 57-67.
- BIGGINS, J.D. (1996) How fast does a general branching random walk spread? "Classical and Modern Branching Processes", *IMA 84, Athreya and Jagers, Eds* Springer.
- BIGGINS, J.D. (1996) Autres
- BIGGINS, J.D. and BINGHAM, N.H. (1993) Large Deviations in the Supercritical Branching Process *Adv. Appl. Prob.* **25** 757-772.
- BRAMSON, M.D. (1978) Maximal displacement of branching Brownian motion *Commun. Pure and Appl. Math.* **31** 531-581.
- BRAMSON, M.D. NEY, P. and TAO, J. (1992) Composition of multitype branching populations *Annals of Applied Probability* **2** 575-596
- CHAUVIN B. (1986) Thèse, Université Paris 6.
- CHAUVIN B. and ROUAULT A. (1996) Boltzmann-Gibbs weights in the branching random walk . "Classical and Modern Branching Processes", *IMA 84, Athreya and Jagers, Eds* Springer.
- CHAUVIN B. and ROUAULT A. (1988) KPP equation and supercritical branching brownian motion in subcritical areas. Application to spatial trees. *Prob.Theory and Related Fields* **79** 299-314.
- CHAUVIN B. and ROUAULT A. (1990) KPP equation and branching brownian motion. The critical case. *Math.Nach.*, **149** 41-59.
- COHEN, S. and ROSSIGNOL, S. (1997) KPP equation in R^d . *Preprint Université de Versailles*.
- DEMBO A, ZEITOUNI O. (1993) *Large deviations techniques and applications*, Jones and Bartlett.
- DEUSCHEL, J.D. and Stroock D.W. (1989) *Large deviations*, Academic Press.
- DION, J.P. (1991) *Statistical inference for discrete time branching processes*. 7th International Summer School on Probability Theory and Mathematical Statistics. Varna. (Obretemov and Stefanov Eds.).
- DUBY, C. and ROUAULT A. (1982) Estimation simultanée de l'espérance et de la variance pour un processus de Galton-Watson. *Ann.IHP.* **18** 149-163.
- DUPUIS, P. and ELLIS R.S. (1997) *A weak convergence approach to the theory of large deviations*, Wiley.
- FREIDLIN, M.I. (1990) *Semi-linear pde's and limit theorems for large deviations* Ecole d'Eté de Probabilités de Saint-Flour XX, *Lect. Notes in Math.* **1527**, Springer.
- HARRIS, T.E. (1948) Branching processes *Ann. Math. Statist.* **41** 474-494.

- IKEDA, N. NAGASAWA, M> and WATANABE S. (1968-1969) *J. Math. Kyoto Univ.* **8** 233-277, 365-410 and **9** 97-162.
- KASAHARA Y. (1978) Tauberian theorems of exponential type *J. Math Kyoto Univ.* **18** 209-219.
- KOLMOGOROV, A., PETROWSKI, I. and PISCOUNOV, N. (1937) Etude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique. *Mosc. Univ. Math. Bull.* **1** 1-25.
- KUCZMA, M., CHOCZEWSKI, B. and GER, R. (1990) *Iterative functional equations*, Cambridge University Press.
- KLEBANER F.C. (1996) Population and Density Dependent Branching Processes. "Classical and Modern Branching Processes", *IMA 84, Athreya and Jagers, Eds* Springer.
- KLEBANER F.C. and ZEITOUNI, O. (1994) The exit problem for a class of density dependent branching systems *Ann. Appl. Probab.* **4** 1188-1205.
- KOLMOGOROV, A. PETROVSKII, I., and PISKUNOV, N. (1937) Etude de l'équation de la diffusion avec une croissance de la matière et son application à un problème biologique. *MOSCOW UNIV. MATH. BULL.* **1** 1-25.
- LAREDO, C. and ROUAULT, A. (1983) Grandes déviations, dynamique de populations et phénomènes malthusiens *Ann. IHP*, **19** 323-350.
- LALLEY, S.P. and SELKE, T. (1988) Travelling waves in inhomogeneous branching Brownian motions. I, *Ann. Probab.*, **16** 1051-1062.
- LALLEY, S.P. and SELKE T. (1989) Travelling waves in inhomogeneous branching Brownian motions. II, *Ann. Probab.* **17** 116-127.
- LALLEY, S.P. and SELKE T. (1997) Hyperbolic branching brownian motion. *Preprint Purdue University.*
- LEE T.Y. (1992) Some large deviation theorems for branching diffusions *Ann. Probab.* **20** 1288-1309.
- LIU, Q. (1996) The growth of an entire characteristic function and the tail probabilities of the limit of a tree martingale *Trees*, Cohen, Chauvin and Rouault Eds. Birkhauser
- LYONS, R. (1997) A simple path to Biggins' martingale convergence in the branching random walk. In *Classical and modern branching processes*, (K.B. Athreya and P. Jagers Editors) IMA Volume 84.
- LYONS, R. (1998) Probability and trees.
- NEVEU, J (1986) Arbres et processus de Galton–Watson *Ann. IHP* **22**, No. 2, 199-207.
- NEVEU, J (1988) Multiplicative martingales for spatial branching processes. In *Seminar Stoch. Proc.* Princeton 223-242, Birkhauser.
- NEY, P. Branching random walk (1992) In *Progress in Probability* **19**, 3-22, Birkhauser.
- PIERRE-LOTI-VIAUD D. and PORTAL F. (1993) Large deviations for random perturbations of discrete time dynamical systems *Bull. Sc. Math.* **117** 333-355.
- ROUAULT A. (1987) Probabilités de présence dans un processus de branchement spatial markovien *Ann.IHP*, **23** 37-61
- ROUAULT A. (1987) Espérances et majorations pour un processus de branchement spatial markovien. *Ann.IHP* **23** 459-497.
- ROUAULT A. (1993) Precise estimates of presence probabilities in the branching random walk. *Stoch.Proc.and their Appl*, **44** 27-39.
- ROUQUES J-PH. (1997) Laplace asymptotics for generalized KPP equation. *ESAIMPS*
- SHWARTZ A. and WEISS, A. (1996) *Large deviations for performance analysis*, Chapman and Hall.
- STROOCK D.W. (1984) *An introduction to the theory of large deviations*, Springer.
- UCHIYAMA, K. (1982) Spatial growth of a branching process of particles living in \mathbb{R}^d *Ann. Probab.* **10** 896-918.
- VARADHAN S.R.S. (1984) *Large deviations and Applications*, SIAM.
- VATUTIN (1993) Branching Processes II *Itogi Nauki i tekhniki* **2** 1-90.
- WAYMIRE (1998) This volume.

ALAIN ROUAULT,
DEPT DE MATHS

UNIVERSITÉ DE VERSAILLES
BAT. FERMAT
45, AVENUE DES ETATS-UNIS
78035 VERSAILLES CEDEX
EMAIL ROUAULT@MATH.UVSQ.FR